

# Lancaster County Department of Community Corrections

## URINE ABSTINENCE TESTING AND INCIDENTAL ALCOHOL EXPOSURE CONTRACT

It is **YOUR** responsibility to limit your exposure to the products and substances detailed below that contain ethyl alcohol. It is **YOUR** responsibility to read product labels, to know what is contained in the products you use and consume and to stop and inspect these products **BEFORE** you use them. **Use of the products detailed below in violation of this contract will NOT be allowed as an excuse for a positive test result. When in doubt, don't use, consume or apply.**

**Cough syrups and other liquid medications:** Participants are prohibited from using alcohol-containing cough/cold syrups, such as Nyquil®. Other cough syrup brands and numerous other liquid medications, rely upon ethyl alcohol as a solvent. Participants are required to read product labels carefully to determine if they contain ethyl alcohol (ethanol). Information on the composition of prescription medications should be available upon request from your pharmacist. Non-alcohol containing cough and cold remedies are readily available at most pharmacies and major retail stores.

**Non-Alcoholic Beer and Wine:** Although legally considered non-alcoholic, non-alcoholic beers (e.g. O'Douls®, Sharps®) do contain a residual amount of alcohol that may result in a positive test for alcohol, if consumed. Participants are **not** permitted to ingest non-alcoholic beer or non-alcoholic wine.

**Food and Other Ingestible Products:** There are numerous other consumable products that contain ethyl alcohol that could result in a positive test for alcohol. Flavoring extracts, such as vanilla or almond extract, and liquid herbal extracts (such as Ginkgo Biloba), could result in a positive screen for alcohol or its breakdown products. Communion wine, food cooked with wine, and flambé dishes (alcohol poured over a food and ignited such as cherries jubilee or baked Alaska) must be avoided. Read carefully the labels on any liquid herbal or homeopathic remedy.

**Mouthwash and Breath Strips:** Most mouthwashes (Listerine®, Cepacol®, etc.) and other breath cleansing products contain ethyl alcohol. The use of mouthwashes containing ethyl alcohol can produce a positive test result. Participants are required to read product labels and educate themselves as to whether a mouthwash product contains ethyl alcohol. Use of ethyl alcohol-containing mouthwashes and breath strips by participants is not permitted. Non-alcohol mouthwashes are readily available and are an acceptable alternative.

**Hand sanitizers:** Hand sanitizers (e.g. Purell®, Germex®, etc.) and other antiseptic gels and foams used to disinfect hands can contain up to 70% ethyl alcohol. Excessive, unnecessary or repeated use of these products could result in a positive urine test. Hand washing with soap and water are just as effective for killing germs.

**Hygiene Products:** Aftershave and colognes, hair sprays and mousse, astringents, insecticides (bug sprays such as Off®) and some body washes contain ethyl alcohol. While it is unlikely that limited use of these products would result in a positive test for alcohol (or its breakdown products), excessive, unnecessary or repeated use of these products could affect test results. Participants must use such products sparingly to avoid reaching detection levels. Just as the court requires participants to regulate their fluid intake to avoid dilute urine samples, it is likewise incumbent upon each participant to limit their use of topically applied (on the skin) products containing ethyl alcohol.

**Solvents and Lacquers.** Many solvents, lacquers and surface preparation products used in industry, construction, and the home contain ethyl alcohol. Both excessive inhalation of vapors, and topical exposure to such products, can potentially cause a positive test result for alcohol. As with the products noted above, participants must educate themselves as to the ingredients in the products they are using. There are alternatives to nearly any item containing ethyl alcohol. Frequency of use and duration of exposure to such products should be kept to a minimum. A positive test result will not be excused by reference to use of an alcohol-based solvent.

### **I HAVE READ AND UNDERSTAND MY RESPONSIBILITIES:**

\_\_\_\_\_  
PARTICIPANT SIGNATURE

\_\_\_\_\_  
PRINTED NAME

\_\_\_\_\_  
DATE